




Fig 1#



09/819,401 - Docket No. 397272000700

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTION


Inventor: Laurent HUMEAU et al

Application No.: 09/819,401 - Docket No. 397272005/00

Sheet 2 of 49




Fig 1B





09819401-031001



Fig 1C




09819401.091001



09819401.093001



Fig 1E



09819401.001001



Fig 1F






Fig 16






Fig 1H





Fig 1 I



09819401-001001

F₁₀ 15.





Fig 1K



09819401-091001



A +105 GTGTGCCCGTCTG +117
B AC . . .

A +118 TTGTGTGACTCTG +130
B

A +131 GTAAGTAGAGATC +143
B . C . G A .


FIG. 2

0919401-091001

FIGURE 13

3A

Ratio Optimization for pN1(cPTC)ASenvGFP Vector






TEOT50"TE046TE60

24

Ratio Optimization for pN1(cPT)GFP Vectors

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUMEAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 14 of 49






FOOTER: FEB 19 2009

3C

Ratio Optimization for pN1(cPT2)ASenvGFP Vector






FOOTNOTES

3b

Best Vector to Packaging Ratio for pN1cGFP Vector

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUMEAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 16 of 49





FOOTED" T046T860

3E

Optimization of vector to packaging ratio for pN2cGFP

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUMEAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 17 of 49





Fig 4A



0931601046T860

Page 4B







FIG. 10

5A


Ratio Optimization for Packaging of pS1cGFP vectors.




TOOT60" T046T860


58

Optimization of vector to packaging ratio for pS2cGFP





Packaging Construct



New features:

- First 42 nt of gag are degenerated.
- Tat and rev represented as cDNA.
- First 208 nt of rev and last 183 nt of tat are degenerated.
- RRE from HIV-2 is used instead of HIV-1 RRE.


These features eliminate almost any homology with the vector plasmid, make system safer.

- Anti-U5 ribozyme is expressed within gag/pol/RRE cassette, further improving safety.
- Gag/pol/rev/tat/RRE cassette and VSV-G expressed from the same plasmid. This feature may enhance packaging efficiency and titers of the vectors.

09/819,401-091001




Fig. 6B Packaging Plasmid
for Second Generation
Vectors



09819401.091001




Fig. 6C Packaging Plasmid
for First Generation Vectors




09819401-031001

Fig 6 D




09/819,401-091001

Fig 6E




09819401.051001

Fig 6F



09819401-091001

Fig 66





09819401-0001

FIG. 1

Fig

Influence of Ribozyme(s) in the Packaging on pN1(cPT)GFP Vector Titers in HeLa-tat Cells






TOOT60" T046T860

Fig 8

Challenge #26, MOI 0.1, 100% transduced






Figure 9A

FOOTNOTES




FIGURE 9B

Potent Inhibition of Wild-type HIV Replication
by Smartvector Containing T Cells





FIG. 10A


Fig 10A



T00T60" T046T860

10/5

Expansion of SupT1 cells after BG & BCNU









Fig 12



Marker

1 pN1 CGFP 1C exp 30

3 pN1 CGFP 2C exp 30

1-4 pVP1.2

9-12 pVP1.2 Rz

13-16 pVP1.2 Rz2

pNL4-3 with DNase I

pNL4-3 without DNase I


Amp. Neg. Control

Extraction Neg. Control

Marker

100T60" T046T860

Fig 13A



Histogram Statistics

File: SupT1 exp.52 transd..009
Tube: pN1(cPT)ASenvGFP 452 a

Sample ID: SupT1 ex
Acquisition Date: 25-

Marker	Left, Right	Events	% Gated	% Total	Mean
All	1, 9910	6356	100.00	63.56	570.39
M1	1, 20	95	1.49	0.95	13.86
M2	20, 9910	6262	98.52	62.62	578.74

0981401-091001

FIGURE 13B
Fig 13B






Fig 14A

Vsv-G, RD114 AND RD114-VSV-G CHIMERIA ENVELOPE PROTEINS





Fig 14B

Titers of RD114-pseudotyped HIV-1 vectors in HT1080

Envelopes	IU/ml
VSV G	3.5x10e6
Rabies virus G	1.6x10e6
RD114WT env	1.5x10e5
RD114E env	3.8x10e4




Fig 15A



09819401.021001




Fig 15B



09819401-091001




Fig 15C



09819401.091001




Fig 15D



09/819,401 - Docket No. 397272000700

Fig 15 E



0501051001

Rev dependent VSV-G constructs




Figure 2



Yield of pN1(cPT)GFP Vectors per Cell Factory before and after Concentration in HeLa-tat Cells.




Fig 16

BEFORE
CONCENTRATION


AFTER
CONCENTRATION





+ = pcuv-Rev
- = PCI
G = β -globin SD
IM-HIV-1 major SD
H-Haymanskydd's SD
IE-HIV-1 env SD
2E-HIV-2 env SD

REMOVE TETRACYCLINE
TO INDUCE EXPRESSION OF VSV-G
THAT IS ~~DEPENDENT~~ DEPENDENT.



TETRACYCLINE	1	2	3	4	5	6	7	8	9	10	11	12	13					
+	+	+	+	+	+	+	+	+	+	+	+	+	+					
-	-	-	-	-	-	-	-	-	-	-	-	-	-					
	X pcuv-vsv-g			pcuv-RRE-4			pcuv-RRE-IM			pcuv-RRE-H			pcuv-RRE-IE			pcuv-RRE-2E		

293G


LANE

001917001046T860

July 17

F.2 18

Influence of the Buffer on Vector Recovery after Storage for 3-5 Weeks at Different Temperatures



- D-PBS
- D-PBS+5% mannitol 1:1
- D-PBS+10% trehalose 1:1
- D-PBS+10% glucose 1:1
- D-PBS+10% trehalose 1:1







Figure 19



09/819,401 - Docket No. 397272000700